

11

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
3 July 2003 (03.07.2003)

PCT

(10) International Publication Number
WO 03/053930 A1

(51) International Patent Classification⁷: C07D 211/90,
401/04, A61K 31/44 // (C07D 401/04, 213:00, 211:00)

(21) International Application Number: PCT/EP02/13931

(22) International Filing Date: 9 December 2002 (09.12.2002)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
0130553.1 20 December 2001 (20.12.2001) GB
0216664.3 17 July 2002 (17.07.2002) GB

CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH,
GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC,
LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW,
MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG,
SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC,
VN, YU, ZA, ZM, ZW.

(84) Designated States (regional): ARIPO patent (GH, GM,
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW),
Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM),
European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE,
ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, SI, SK,
TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ,
GW, ML, MR, NE, SN, TD, TG).

(71) Applicant (for all designated States except US): BAYER
AKTIENGESELLSCHAFT [DE/DE]; 51368 Lev-
erkusen (DE).

(72) Inventors; and

(75) Inventors/Applicants (for US only): GIELEN, Heike
[DE/DE]; Am Kettnersbusch 3, 51379 Leverkusen (DE).
LI, Volkhart, Min-Jian [DE/DE]; Im Wiesengrund
40, 42553 Velbert (DE). ROSENRETER, Ulrich
[DE/DE]; Obere Rutenbeck 6, 42349 Wuppertal (DE).
SCHLEMMER, Karl-Heinz [DE/DE]; Wildsteig 22
a, 42113 Wuppertal (DE). ALLERHEILIGEN, Swen
[DE/DE]; Beckumsfeld 4, 45259 Essen (DE). NASH,
Kevin [GB/GB]; 136 a Littlebrook Avenue, Burnham,
Buckinghamshire SL2 2NE (GB). FITZGERALD, Mary
[GB/GB]; 2 Paternoster Court, Cassington, Yarnton,
Oxfordshire OX5 1QB (GB).

Declaration under Rule 4.17:

— as to applicant's entitlement to apply for and be granted
a patent (Rule 4.17(ii)) for the following designations AE,
AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA,
CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES,
FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE,
KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG,
MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU,
SD, SE, SG, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, UZ,
VC, VN, YU, ZA, ZM, ZW, ARIPO patent (GH, GM, KE, LS,
MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent
(AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent
(AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB,
GR, IE, IT, LU, MC, NL, PT, SE, SI, SK, TR), OAPI patent
(BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE,
SN, TD, TG)

Published:

— with international search report

(74) Common Representative: BAYER AKTIENGE-
SELLSCHAFT; 51368 Leverkusen (DE).

(81) Designated States (national): AE, AG, AL, AM, AT, AU,
AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU,

For two-letter codes and other abbreviations, refer to the "Guid-
ance Notes on Codes and Abbreviations" appearing at the begin-
ning of each regular issue of the PCT Gazette.

(54) Title: 1,4-DIHYDRO-1,4-DIPHENYLPYRIDINE DERIVATIVES

(57) Abstract: The invention relates to novel 1,4-dihydro-1,4-diphenylpyridine derivatives, processes for their preparation, and their use in medicaments, especially for the treatment of chronic obstructive pulmonary diseases.



WO 03/053930 A1

1,4-Dihydro-1,4-diphenylpyridine derivatives


The present invention relates to novel 1,4-dihydro-1,4-diphenylpyridine derivatives, processes for their preparation, and their use in medicaments, especially for the treatment of chronic obstructive pulmonary diseases.

The fibrous protein elastin, which comprises an appreciable percentage of all protein content in some tissues, such as the arteries, some ligaments and the lungs, can be hydrolysed or otherwise destroyed by a select group of enzymes classified as elastases. Human leukocyte elastase (HLE, EC 3.4.21.37), also known as human neutrophil elastase (HNE), is a glycosylated, strongly basic serine protease and is found in the azurophilic granules of human polymorphonuclear leukocytes (PMN). HNE is released from activated PMN and has been implicated causally in the pathogenesis of acute and chronic inflammatory diseases. HNE is capable of degrading a wide range of matrix proteins including elastin, and in addition to these actions on connective tissue HNE has a broad range of inflammatory actions including upregulation of IL-8 gene expression, oedema formation, mucus gland hyperplasia and mucus hypersecretion. Pulmonary diseases where HNE is believed to play a role include lung fibrosis, pneumonia, acute respiratory distress syndrome (ARDS), pulmonary emphysema, including smoking-induced emphysema, chronic obstructive pulmonary diseases (COPD) and cystic fibrosis. HNE has also been causally implicated in rheumatoid arthritis, atherosclerosis, brain trauma, cancer and related conditions in which neutrophil participation is involved. Thus, inhibitors of HLE activity can be potentially useful in the treatment of a number of inflammatory diseases, especially of chronic obstructive pulmonary diseases [R.A. Stockley, *Neutrophils and protease/antiprotease imbalance*, Am. J. Respir. Crit. Care 160, S49-S52 (1999)].

Certain cationic amphiphilic 1,4-dihydropyridine derivatives useful for delivery of nucleotide containing compounds are disclosed in WO-A1-01/62946. Ethyl 6-amino-1,4-bis(4-chlorophenyl)-5-cyano-2-methyl-1,4-dihydro-3-pyridinecarboxylate has been

synthesized and tested for potential antimicrobial activity as described in A.W. Erian et al., *Pharmazie* 53 (11), 748-751 (1998).

The present invention relates to compounds of the general formula (I)



5

wherein

R^1 , R^2 , R^3 , R^8 , R^9 and R^{10} independently from each other represent hydrogen, halogen, nitro, cyano, trifluoromethyl, C_1 - C_6 -alkyl, C_1 - C_6 -alkanoyl, hydroxy, C_1 - C_6 -alkoxy, trifluoromethoxy, amino, mono- or di- C_1 - C_6 -alkylamino, C_1 - C_6 -acylamino, C_1 - C_6 -alkoxycarbonylamino, carboxyl, C_1 - C_6 -alkoxy-carbonyl or phenyl, wherein C_1 - C_6 -alkyl, C_1 - C_6 -alkanoyl, C_1 - C_6 -alkoxy, mono- or di- C_1 - C_6 -alkylamino and C_1 - C_6 -acylamino can be further substituted with one to three identical or different radicals selected from the group consisting of hydroxy, C_1 - C_4 -alkoxy, amino, mono- and di- C_1 - C_4 -alkylamino,

15

R^4 represents C_1 - C_6 -alkyl, trifluoromethyl or phenyl,

R^5 represents C_1 - C_4 -alkyl, which can be substituted with one to three identical or different radicals selected from the group consisting of hydroxy, C_1 - C_4 -alkoxy, amino, mono- and di- C_1 - C_4 -alkylamino,

20


- 3 -

R⁶ represents cyano, aminocarbonyl, mono- and di-C₁-C₄-alkylaminocarbonyl, carboxyl or C₁-C₆-alkoxycarbonyl, wherein the alkoxy moiety can be further substituted with a radical selected from the group consisting of hydroxy, C₁-C₄-alkoxy, amino, mono- and di-C₁-C₄-alkylamino,

5

or

R⁶ represents a moiety of the formula



10

or

wherein R^A is selected from the group consisting of hydrogen and C₁-C₆-alkyl,

15

R⁷ represents hydrogen, C₁-C₄-alkyl or amino,

X¹, X², X³ and X⁴ independently from each other represent CH or N, wherein ring A contains either 0, 1 or 2 nitrogen atoms,

20

and

Y¹, Y², Y³, Y⁴ and Y⁵ independently from each other represent CH or N, wherein ring B contains either 0, 1 or 2 nitrogen atoms.

25

In another embodiment, the present invention relates to compounds of general formula (I), wherein

- 4 -

R¹ represents hydrogen,

5 R², R³, R⁹ and R¹⁰ independently from each other represent hydrogen, halogen, nitro, cyano, trifluoromethyl, C₁-C₄-alkyl, C₁-C₄-alkanoyl, hydroxy, C₁-C₄-alkoxy, trifluoromethoxy, amino, C₁-C₄-dialkylamino, C₁-C₄-acylamino, methoxy-carbonylamino, tert.-butoxycarbonylamino, carboxyl, methoxycarbonyl or ethoxycarbonyl, wherein C₁-C₄-alkyl, C₁-C₄-alkanoyl, C₁-C₄-alkoxy, C₁-C₄-dialkylamino and C₁-C₄-acylamino can be further substituted with one to two
10 identical or different radicals selected from the group consisting of hydroxy, methoxy, ethoxy, amino, dimethylamino and diethylamino,

R⁴ represents methyl, ethyl, trifluoromethyl or phenyl,

15 R⁵ represents methyl or ethyl,

R⁶ represents cyano, aminocarbonyl, methylaminocarbonyl, dimethylaminocarbonyl, carboxyl or C₁-C₄-alkoxycarbonyl, wherein the alkoxy moiety can be further substituted with a radical selected from the group consisting of hydroxy, methoxy, ethoxy, amino, mono- and di-C₁-C₄-alkylamino,
20

R⁷ represents hydrogen, methyl, ethyl or amino,

R⁸ represents hydrogen,

25 X¹, X² and X³ represent CH,

X⁴ represents CH or N,

and

30

Y¹, Y², Y³, Y⁴ and Y⁵ represent CH.

In another embodiment, the present invention relates to compounds of general formula (I), wherein

5 R^1 and R^2 represent hydrogen,

R^3 represents fluoro, chloro, bromo, nitro, cyano, trifluoromethyl, methyl, methoxy or hydroxy,

10 R^4 represents methyl or trifluoromethyl,

R^5 represents methyl,

15 R^6 represents cyano, aminocarbonyl, methylaminocarbonyl, dimethylaminocarbonyl, carboxyl, methoxycarbonyl or ethoxycarbonyl,

R^7 represents hydrogen, methyl or amino,

20 R^8 and R^9 represent hydrogen,

R^{10} represents fluoro, chloro, bromo, nitro, cyano, trifluoromethyl or methyl,


X^1 , X^2 , X^3 and X^4 represent CH,

25 and

Y^1 , Y^2 , Y^3 , Y^4 and Y^5 represent CH.

30 In another embodiment, the present invention relates to compounds according to general formula (I), which have the following structure:

- 6 -



and wherein

5 R^1, R^2, R^3, R^8, R^9 and R^{10} independently from each other represent hydrogen, halogen, nitro, cyano, trifluoromethyl, C_1 - C_6 -alkyl, C_1 - C_6 -alkanoyl, hydroxy, C_1 - C_6 -alkoxy, trifluoromethoxy, amino, mono- or di- C_1 - C_6 -alkylamino, C_1 - C_6 -acylamino, C_1 - C_6 -alkoxycarbonylamino, carboxyl, C_1 - C_6 -alkoxycarbonyl or phenyl, wherein C_1 - C_6 -alkyl, C_1 - C_6 -alkanoyl, C_1 - C_6 -alkoxy, mono- or di- C_1 - C_6 -alkylamino and C_1 - C_6 -acylamino can be further substituted with one to three identical or
 10 different radicals selected from the group consisting of hydroxy, C_1 - C_4 -alkoxy, amino, mono- and di- C_1 - C_4 -alkylamino,

15 R^4 represents C_1 - C_6 -alkyl, trifluoromethyl or phenyl,

R^5 represents C_1 - C_4 -alkyl,

20 R^6 represents cyano, carboxyl or C_1 - C_6 -alkoxycarbonyl, wherein the alkoxy moiety can be further substituted with a radical selected from the group consisting of hydroxy, C_1 - C_4 -alkoxy, amino, mono- and di- C_1 - C_4 -alkylamino,

and

R^7 represents hydrogen, C_1 - C_4 -alkyl or amino.

5 In another embodiment, the present invention relates to compounds according to general formula (I), wherein R^3 is cyano, which is located in para-position relativ to the 1,4-dihydropyridine ring.

In another embodiment, the present invention relates to compounds according to
10 general formula (I), wherein R^4 is methyl.

In another embodiment, the present invention relates to compounds according to general formula (I), wherein R^5 is methyl.

15 In another embodiment, the present invention relates to compounds according to general formula (I), wherein R^6 is methoxycarbonyl or ethoxycarbonyl.

In another embodiment, the present invention relates to compounds according to general formula (I), wherein R^7 is hydrogen, methyl or amino.
20

In another embodiment, the present invention relates to compounds according to general formula (I), wherein R^{10} is trifluoromethyl, which is attached to Y^2 .

The compounds according to this invention can also be present in the form of their
25 salts, hydrates and/or solvates.

Physiologically acceptable salts are preferred in the context of the present invention.

Physiologically acceptable salts according to the invention are non-toxic salts which
30 in general are accessible by reaction of the compounds (I) with an inorganic or organic base or acid conventionally used for this purpose. Non-limiting examples of

pharmaceutically acceptable salts of compounds (I) include the alkali metal salts, e.g. lithium, potassium and sodium salts, the alkaline earth metal salts such as magnesium and calcium salts, the quaternary ammonium salts such as, for example, triethyl ammonium salts, acetates, benzene sulphonates, benzoates, dicarbonates, disulphates, 5 ditartrates, borates, bromides, carbonates, chlorides, citrates, dihydrochlorides, fumarates, gluconates, glutamates, hexyl resorcinates, hydrobromides, hydrochlorides, hydroxynaphthoates, iodides, isothionates, lactates, laurates, malates, maleates, mandelates, mesylates, methylbromides, methylnitrates, methylsulphates, nitrates, oleates, oxalates, palmitates, pantothenates, phosphates, diphosphates, 10 polygalacturonates, salicylates, stearates, sulphates, succinates, tartrates, tosylates, valerates, and other salts used for medicinal purposes.

Hydrates of the compounds of the invention or their salts are stoichiometric compositions of the compounds with water, such as for example hemi-, mono-, or 15 dihydrates.

Solvates of the compounds of the invention or their salts are stoichiometric compositions of the compounds with solvents.

20 The present invention includes both the individual enantiomers or diastereomers and the corresponding racemates or diastereomeric mixtures of the compounds according to the invention and their respective salts. In addition, all possible tautomeric forms of the compounds described above are included according to the present invention. The diastereomeric mixtures can be separated into the individual isomers by 25 chromatographic processes. The racemates can be resolved into the respective enantiomers either by chromatographic processes on chiral phases or by resolution.

In the context of the present invention, the substituents, if not stated otherwise, in general have the following meaning:

Alkyl in general represents a straight-chain or branched hydrocarbon radical having 1 to 6, preferably 1 to 4 carbon atoms. Non-limiting examples include methyl, ethyl, n-propyl, isopropyl, n-butyl, isobutyl, sec.-butyl, tert.-butyl, pentyl, isopentyl, hexyl, isohexyl. The same applies to radicals such as alkoxy, alkylamino, alkoxy carbonyl and alkoxy carbonylamino.

Acyl or Alkanoyl in general represents a straight-chain or branched hydrocarbon radical having 1 to 6, preferably 1 to 4 carbon atoms which has a carbonyl function at the position of attachment. Non-limiting examples include formyl, acetyl, n-propionyl, n-butyryl, isobutyryl, pivaloyl, n-hexanoyl. The same applies to radicals such as acylamino.

Alkylamino represents an alkylamino radical having one or two (independently selected) alkyl substituents, illustratively and preferably representing methylamino, ethylamino, n-propylamino, isopropylamino, tert-butylamino, n-pentylamino, n-hexylamino, *N,N*-dimethylamino, *N,N*-diethylamino, *N*-ethyl-*N*-methylamino, *N*-methyl-*N*-n-propylamino, *N*-isopropyl-*N*-n-propylamino, *N*-t-butyl-*N*-methylamino, *N*-ethyl-*N*-n-pentylamino and *N*-n-hexyl-*N*-methylamino.

Alkylaminocarbonyl represents an alkylaminocarbonyl radical having one or two (independently selected) alkyl substituents, illustratively and preferably representing methylaminocarbonyl, ethylaminocarbonyl, n-propylaminocarbonyl, isopropylaminocarbonyl, tert-butylaminocarbonyl, n-pentylaminocarbonyl, n-hexylaminocarbonyl, *N,N*-dimethylaminocarbonyl, *N,N*-diethylaminocarbonyl, *N*-ethyl-*N*-methylaminocarbonyl, *N*-methyl-*N*-n-propylaminocarbonyl, *N*-isopropyl-*N*-n-propylaminocarbonyl, *N*-t-butyl-*N*-methylaminocarbonyl, *N*-ethyl-*N*-n-pentylamino-carbonyl and *N*-n-hexyl-*N*-methylaminocarbonyl.

When stated, that X^1 , X^2 , X^3 and X^4 as well as Y^1 , Y^2 , Y^3 , Y^4 and Y^5 represent CH or N, CH shall also stand for a ring carbon atom, which is substituted with a substituent R^1 , R^2 , R^3 or R^8 , R^9 and R^{10} , respectively.


- 10 -

A * symbol next to a bond denotes the point of attachment in the molecule.

The letters A and B are used to designate the different rings and, for the purpose of clarity, are not shown in all structures.



In another embodiment, the present invention relates to processes for synthesizing the compounds of general formula (I), characterized in that either

10 [A] compounds of the general formula (II)



wherein R^4 , R^5 , R^8 , R^9 , R^{10} and Y^1 to Y^5 have the meaning described above,

15 are condensed in the presence of a base, in a three-component-reaction, with compounds of the general formulas (III) and (IV)




wherein R^1 , R^2 , R^3 and X^1 to X^4 have the meaning described above,

20

and

R^{11} represents cyano or C_1 - C_6 -alkoxycarbonyl,


to give compounds of the general formula (Ia)



5 or


[B] compounds of the general formulas (II) and (III) are condensed in the presence of an acid, in a three-component-reaction, with compounds of the general formula (V)

10



wherein R^{12} represents C_1 - C_6 -alkyl,


15 to give compounds of the general formula (Ib)



or

[C] compounds of the general formula (VI)

5




wherein R⁸, R⁹, R¹⁰, R¹² and Y¹ to Y⁵ have the meaning described above,

10 and

R¹³ represents C₁-C₄-alkyl,


are condensed in the presence of an acid or a base, in a three-component-reaction, with compounds of the general formulas (III) and (VII)



wherein R^4 and R^5 have the meaning described above,


to give compounds of the general formula (Ic)

5



The processes can be illustrated by the following schemes [A] to [C]:

Scheme [A] to [C]



Suitable solvents for the processes [A] to [C] are generally customary organic solvents which do not change under the reaction conditions. These include ethers such as diethyl ether, diisopropyl ether, 1,2-dimethoxyethane, dioxan or tetrahydrofuran, ethylacetate, acetone, acetonitrile, dimethylsulfoxide, dimethylformamide, or alcohols such as methanol, ethanol, n-propanol, isopropanol, n-butanol or t-butanol, or hydrocarbons such as pentane, hexane, cyclohexane, benzene, toluene or xylene, or halogeno-hydro-

carbons such as dichloromethane, dichloroethane, trichloromethane or chlorobenzene. For process [B] and [C] also acetic acid can be employed as solvent. It is also possible to use mixtures of the above-mentioned solvents. Preferred for process [A] is ethanol or a mixture of n-propanol and dimethylsulfoxide. For process [B] and [C] acetic acid or diisopropyl ether are preferred.

Suitable bases for process [A] and [C] are generally inorganic or organic bases. These preferably include cyclic amines, such as, for example, piperidine, morpholine, N-methylmorpholine, pyridine or 4-N,N-dimethylaminopyridine, or (C₁-C₄)-trialkylamines, such as, for example, triethylamine or diisopropylethylamine. Preference is given to piperidine. The base is employed in an amount from 0.1 mol to 10 mol, preferably from 1 mol to 3 mol, relative to 1 mol of the compound of the general formula (II).


Suitable acids for process [B] and [C] are generally inorganic or organic acids. These preferably include carboxylic acids, such as, for example acetic acid or trifluoroacetic acid, or sulfonic acids, such as, for example, methanesulfonic acid or p-toluenesulfonic acid. Preference for process [B] and [C] is given to acetic acid or trifluoroacetic acid. The acid is employed in an amount from 0.25 mol to 100 mol, relative to 1 mol of the compounds of the general formulas (V) and (VII), respectively.

The processes [A] to [C] are in general carried out in a temperature range from +20°C to +150°C, preferably from +60°C to +130°C.

The processes [A] to [C] are generally carried out at normal pressure. However, it is also possible to carry them out at elevated pressure or at reduced pressure (for example in a range from 0.5 to 5 bar).

The compounds of general formula (II) can be synthesized by reacting compounds of general formula (VII) with compounds of the general (VIII)

- 16 -



wherein R^8 , R^9 , R^{10} and Y^1 to Y^5 have the meaning described above,

- 5 in the presence of an acid, such as acetic acid or p-toluenesulfonic acid, under water-removing conditions.

The compounds of general formula (VI) can be synthesized analogously.

- 10 The compounds of the general formulas (III), (IV), (V), (VII) and (VIII) are known per se, or they can be prepared by customary methods.

Surprisingly, the compounds of the present invention show human neutrophil elastase (HNE) inhibitory activity and are therefore suitable for the preparation of medicaments for the treatment of diseases associated with HNE activity. They may thus provide an effective treatment of acute and chronic inflammatory processes, such as rheumatoid arthritis, atherosclerosis, and especially of acute and chronic pulmonary diseases, such as lung fibrosis, cystic fibrosis, pneumonia, acute respiratory distress syndrome (ARDS), in particular pulmonary emphysema, including smoking-induced emphysema, and chronic obstructive pulmonary diseases (COPD). They may also provide an effective treatment of brain trauma, cancer and other conditions in which neutrophil participation is involved.

The compounds of formula (I) according to the invention can therefore be used as active compound components for the production of medicaments. For this, they can be converted into the customary formulations such as tablets, coated tablets, aerosols, pills, granules, syrups, emulsions, suspensions and solutions in a known manner using inert, non-toxic, pharmaceutically suitable excipients or solvents. Preferably,

the compounds according to the invention are used here in an amount such that their concentration in the total mixture is approximately 0.5% to approximately 90% by weight, the concentration, inter alia, being dependent on the corresponding indication of the medicament.

5

The above mentioned formulations are produced, for example, by extending the active compounds with solvents and/or excipients having the above properties, where, if appropriate, additionally emulsifiers or dispersants and, in the case of water as the solvent, alternatively an organic solvent, have to be added.

10

Administration is carried out in a customary manner, preferably orally, transdermally or parenterally, for example perlingually, buccally, intravenously, nasally, rectally or by inhalation.

15

For human use, in the case of oral administration, it is recommendable to administer doses of from 0.001 to 50 mg/kg, preferably of 0.01 mg/kg to 20 mg/kg. In the case of parenteral administration, such as, for example, intravenously or via mucous membranes nasally, buccally or inhalationally, it is recommendable to use doses of 0.001 mg/kg to 0.5 mg/kg.

20

In spite of this, if appropriate, it may be necessary to depart from the amounts mentioned above, namely depending on the body weight or the type of administration route, on the individual response towards the medicament, the manner of its formulation and the time or interval at which administration takes place. Thus, in some cases it may be sufficient to manage with less than the above mentioned minimum amount, while in other cases the upper limit mentioned must be exceeded. In the case of the administration of relatively large amounts, it may be recommendable to divide these into several individual doses over the course of the day.

25

A. Evaluation of physiological activity

The potential of the compounds of the invention to inhibit neutrophil elastase activity may be demonstrated, for example, using the following assays:

I. In vitro assays of human neutrophil elastase (HNE)**Assay contents**

assay buffer: 0.1 M HEPES-NaOH buffer pH 7.4, 0.5 M NaCl, 0.1% (w/v) bovine
serum albumin;
suitable concentration (see below) of HNE (18 U/mg lyophil., #20927.01, SERVA
Electrophoresis GmbH, Heidelberg, Germany) in assay buffer;
suitable concentration (see below) of substrate in assay buffer;
suitable concentration of test compounds diluted with assay buffer from a 10 mM
stock solution in DMSO.

Example A

***In vitro* inhibition of HNE using a fluorogenic peptide substrate (continuous
read-out signal, 384 MTP assay format):**

In this protocol, the elastase substrate MeOSuc-Ala-Ala-Pro-Val-AMC (#324740,
Calbiochem-Novabiochem Corporation, Merck KGaA, Darmstadt, Germany) is used.
The test solution is prepared by mixing 10 µl of test compound dilution, 20 µl of
HNE enzyme dilution (final concentration 8 - 0.4 µU/ml, routinely 2.1 µU/ml) and
20 µl of substrate dilution (final concentration 1 mM - 1 µM, routinely 20 µM),
respectively. The solution is incubated for 0 - 2 hrs at 37°C (routinely one hour). The
fluorescence of the liberated AMC due to the enzymatic reaction is measured at 37°C
(TECAN spectra fluor plus plate reader). The rate of increase of the fluorescence (ex.
395 nm, em. 460 nm) is proportional to elastase activity. IC₅₀ values are determined

by RFU-versus-[I] plots. K_m and $K_{m(app.)}$ values are determined by Lineweaver-Burk plots and converted to K_i values by Dixon plots.

5 The preparation examples had IC_{50} values within the range of 20 nM - 20 μ M in this assay. Representative data are given in table 1:

Example No.	IC_{50} [nM]
3	200
4	40
5	20
8	330
10	230
12	180
13	30
14	20
15	120
17	9000
18	30

table 1

Example B

10

***In vitro* inhibition of HNE using a fluorogenic, unsoluble elastin substrate (discontinuous read-out signal, 96 MTP assay format)**

15 In this protocol the elastase substrate elastin-fluorescein (#100620, ICN Biomedicals GmbH, Eschwege, Germany) is used. The test solution is prepared by mixing 3 μ l of test compound dilution, 77 μ l of HNE enzyme dilution (final concentration 0.22 U/ml - 2.2 mU/ml, routinely 21.7 μ U/ml) and 80 μ l substrate suspension (final concentration 2 mg/ml). The suspension is incubated for 0 - 16 hrs at 37°C (routinely four hours) under slightly shaking conditions. To stop the enzymatic reaction, 160 μ l
20 of 0.1 M acetic acid are added to the test solution (final concentration 50 mM). The

polymeric elastin-fluorescein is pulled down by centrifugation (Eppendorf 5804 centrifuge, 3.000 rpm, 10 min). The supernatant is transferred into a new MTP and the fluorescence of the liberated peptide fluorescein due to the enzymatic reaction is measured (BMG Fluostar plate reader). The rate of fluorescence (ex. 490 nm, em. 520 nm) is proportional to elastase activity. IC₅₀ values are determined by RFU-versus-[I] plots.

II. In vitro PMN elastolysis assay

This assay is used to determine the elastolytic potential of human polymorphonuclear cells (PMNs) and assess the proportion of degradation due to neutrophil elastase [cf. Z.W. She et al., Am. J. Respir. Cell. Mol. Biol. 9, 386-392 (1993)].

Tritiated elastin, in suspension, is coated on to a 96 well plate at 10 µg per well. Test and reference [ZD-0892 (J. Med. Chem. 40, 1876-1885, 3173-3181 (1997), WO 95/21855) and α1 protease inhibitor (α1PI)] compounds are added to the wells at the appropriate concentrations. Human PMNs are separated from peripheral venous blood of healthy donors and resuspended in culture media. The neutrophils are added to the coated wells at concentrations ranging between 1 x 10⁶ to 1 x 10⁵ cells per well. Porcine pancreatic elastase (1.3 µM) is used as a positive control for the assay, and α1PI (1.2 µM) is used as the positive inhibitor of neutrophil elastase. The cellular control is PMNs without compound at each appropriate cell density. The cells plus compounds are incubated in a humidified incubator at 37°C for 4 hours. The plates are centrifuged to allow the harvest of cell supernatant only. The supernatant is transferred in 75 µl volumes to corresponding wells of a 96 well Lumaplate™ (solid scintillant containing plates). The plates are dried until no liquid is visible in the wells and read in a beta counter for 3 minutes per well.

Elastolysis of the ³H-elastin results in an increase in counts in the supernatant. An inhibition of this elastolysis shows a decrease, from the cellular control, of tritium in the supernatant. α1PI gave 83.46 ± 3.97% (mean ± s.e.m.) inhibition at 1.2 µM (n =

3 different donors at 3.6×10^5 cells per well). IC_{50} values are obtained for the reference compound ZD-0892 of 45.50 ± 7.75 nM (mean \pm s.e.m.) (n = 2 different donors at 3.6×10^5 cells per well).

5 Given that ZD-0892 is a selective inhibitor of PMN elastase along with the data from $\alpha 1PI$ inhibition, these results indicate that the majority of elastin degradation by PMNs is due to the release of neutrophil elastase, and not to another elastolytic enzyme such as matrix metalloproteases (MMPs). The compounds of this invention are evaluated for their inhibitory activity in this HNE-dependent model of neutro-phil
10 elastolysis.

III. In vivo model of acute lung injury in the rat

Instillation of human neutrophil elastase (HNE) into rat lung causes acute lung
15 damage. The extent of this injury can be assessed by measuring lung haemorrhage.

Rats are anaesthetised with Hypnorm/Hypnovel/water and instilled with HNE or saline delivered by microsyringe into the lungs. Test compounds are administered by intravenous injection, by oral gavage or by inhalation at set times prior to the
20 administration of HNE. Sixty minutes after the administration of elastase animals are killed by an anaesthetic overdose (sodium pentobarbitone) and the lungs lavaged with 2 ml heparinised phosphate buffered saline (PBS). Bronchoalveolar lavage (BAL) volume is recorded and the samples kept on ice. Each BAL sample is centrifuged at 900 r.p.m. for 10 minutes at $4-10^{\circ}C$. The supernatant is discarded and
25 the cell pellet resuspended in PBS and the sample spun down again. The supernatant is again discarded and the cell pellet resuspended in 1 ml 0.1% cetyltrimethyl-ammonium bromide (CTAB) / PBS to lyse the cells. Samples are frozen until blood content is assayed. Prior to the haemorrhage assay the samples are defrosted and mixed. 100 μ l of each sample are placed into a separate well of a 96 well flat-bottomed plate. All samples are tested in duplicate. 100 μ l 0.1% CTAB/PBS is
30 included as a blank. The absorbance of the well contents is measured at 415 nm using

a spectrophotometer. A standard curve is constructed by measuring the OD at 415 nm of different concentrations of blood in 0.1% CTAB/PBS. Blood content values are calculated by comparison to the standard curve (included in each plate) and normalised for the volume of BAL fluid retrieved.

5

The compounds of this invention are evaluated intravenously, orally or by inhalation for their inhibitory activity in this model of HNE-induced haemorrhage in the rat.

B. Examples**Abbreviations:**

	c	=	concentration
5	DMSO	=	dimethylsulfoxide
	HPLC	=	high performance liquid chromatography
	LC-MS	=	liquid chromatography-coupled mass spectroscopy
	NMR	=	nuclear magnetic resonance spectroscopy
	tlc	=	thin layer chromatography

10

LC-MS Method:

solvent A: acetonitrile

solvent B: 0.3 g 30% HCl/L water


15 column oven: 50°C

column: Symmetry C18 2.1 x 150 mm

gradient :	time [min]	%A	%B	flow [ml/min]
	0	10	90	0.9
	3	90	10	1.2
20	6	90	10	1.2

Starting Materials:25 **Example 1A**

4-[(3-Nitrophenyl)amino]-3-penten-2-one




36.24 g (362 mmol) Acetylacetone, 10.00 g (72 mmol) 3-nitroaniline, and 1.25 g (7.2 mmol) 4-toluenesulfonic acid are dissolved in 100 ml toluene. The reaction mixture is refluxed overnight with a Dean-Stark trap to remove water. After cooling down to room temperature, the solvent is removed *in vacuo* and the residue is purified by column chromatography on silica with eluent dichloromethane.

Yield: 12.0 g (75%)

¹H-NMR (400 MHz, DMSO): δ = 2.0 (s, 3H); 2.1 (s, 3H); 5.4 (s, 1H); 7.7 (m, 2H); 8.0 (m, 2H); 12.5 (s, 1H) ppm.

10 Example 2A

4-[[3-(Trifluoromethyl)phenyl]amino]-3-penten-2-one




15 15.53 g (155 mmol) Acetylacetone, 5.00 g (31 mmol) 3-trifluoromethylaniline, and 0.53 g (3.1 mmol) 4-toluenesulfonic acid are dissolved in 50 ml toluene. The reaction mixture is refluxed overnight with a Dean-Stark trap to remove water. After cooling down to room temperature, the solvent is removed *in vacuo* and the residue is purified by column chromatography on silica with cyclohexane/ethylacetate mixtures as eluent.

Yield: 5.46 g (72%)

¹H-NMR (200 MHz, DMSO): δ = 2.0 (s, 3H); 2.1 (s, 3H); 5.3 (s, 1H); 7.5 (m, 4H); 12.5 (s, 1H) ppm.

Example 3A

Ethyl (2E)-3-[[3-(trifluoromethyl)phenyl]amino]-2-butenolate



5

4.0 g (31 mmol) Ethyl 3-oxobutanoate, 5.0 g (31 mmol) 3-trifluoromethylaniline, and 1.86 g (31 mmol) acetic acid are dissolved in 50 ml toluene. The reaction mixture is refluxed overnight with a Dean-Stark trap to remove water. After cooling down to room temperature, the solvent is removed *in vacuo* and the residue is purified by column chromatography on silica with cyclohexane/ethylacetate mixtures as eluent.

10


Yield: 2.28 g (27%)

¹H-NMR (300 MHz, DMSO): δ = 1.2 (t, 3H); 2.0 (s, 3H); 4.1 (q, 2H); 4.8 (s, 1H); 7.5 (m, 4H); 10.4 (s, 1H) ppm.

15

Example 4A

4-[[3-(Methyl)phenyl]amino]-3-penten-2-one



20

23.36 g (233 mmol) Acetylacetone, 5.00 g (47 mmol) 3-methylaniline, and 0.8 g (4.7 mmol) 4-toluenesulfonic acid are dissolved in 50 ml toluene. The reaction mixture is refluxed overnight with a Dean-Stark trap to remove water. After cooling down to room temperature, the solvent is removed *in vacuo* and the residue is purified by column chromatography on silica with cyclohexane/ethylacetate mixtures as eluent.

25


- 26 -

Yield: 7.7 g (87%)

¹H-NMR (300 MHz, DMSO): δ = 2.0 (2 s, 6H); 2.3 (s, 3H); 5.2 (s, 1H); 7.0 (m, 3H); 7.2 (m, 1H) ppm.

5 **Example 5A**

4-{{3-Iodo-5-(trifluoromethyl)phenyl}amino}-3-penten-2-one




10 1.38 g (13.8 mmol) Acetylacetone, 3.95 g (13.8 mmol) 3-iodo-5-trifluoromethyl-aniline, and 0.1 g (1.6 mmol) 4-toluenesulfonic acid are dissolved in 150 ml toluene. The reaction mixture is refluxed for 7 h with a Dean-Stark trap to remove water. After cooling down to room temperature, the suspension is filtered. The solid is purified by recrystallisation from ethanol.

15 Yield: 1.8 g (34%)

¹H-NMR (300 MHz, DMSO): δ = 2.0 (s, 3H); 2.1 (s, 3H); 5.3 (s, 1H); 7.6 (s, 1H); 7.8 (s, 1H); 7.9 (s, 1H); 12.4 (s, 1H) ppm.

Example 6A

20 4-({3-[(Diethylamino)methyl]phenyl}amino)-3-penten-2-one



25 1.00 g (9.99 mmol) Acetylacetone, 1.78 g (9.99 mmol) 3-[(diethylamino)methyl]-aniline, and 0.15 g (2.5 mmol) 4-toluenesulfonic acid are dissolved in 100 ml

- 27 -


toluene. The reaction mixture is refluxed overnight with a Dean-Stark trap to remove water. 0.05 g (0.83 mmol) 4-toluenesulfonic acid are again added and the mixture is refluxed overnight. As tlc control showed that there is some 3-[(diethylamino)-methyl]aniline left, 0.2 g (2 mmol) acetylacetone are added and refluxing is continued for 4 h. To drive the reaction to completion, 0.6 ml (9.98 mmol) acetic acid and 0.2 g (2 mmol) acetylacetone are added and the mixture is refluxed overnight with a Dean-Stark trap. After cooling down to room temperature, the solvent is removed *in vacuo* and the residue is purified by column chromatography on silica with dichloromethane/methanol mixtures as eluent.

Yield: 1.97 g (76%)

$^1\text{H-NMR}$ (200 MHz, CDCl_3): δ = 1.0 (t, 6H); 2.0 (s, 3H), 2.1 (s, 3H); 2.5 (q, 4H); 3.5 (s, 2H); 5.2 (s, 1H); 7.0 (m, 1H); 7.1-7.3 (m, 3H); 12.5 (s, 1H) ppm.

Example 7A

4-[[3-Methoxy-5-(trifluoromethyl)phenyl]amino]-3-penten-2-one




1.57 g (15.7 mmol) Acetylacetone, 3.00 g (15.7 mmol) 3-methoxy-5-(trifluoromethyl)aniline, and 0.1 g (1.6 mmol) 4-toluenesulfonic acid are dissolved in 150 ml toluene. The reaction mixture is refluxed for 7 h with a Dean-Stark trap to remove water. After cooling down to room temperature, the suspension is filtered. The solid is purified by recrystallisation from ethanol.

Yield: 2.8 g (60%)

$^1\text{H-NMR}$ (300 MHz, DMSO): δ = 2.0 (s, 3H); 2.1 (s, 3H); 3.8 (s, 3H); 5.3 (s, 1H); 7.1 (m, 3H); 12.4 (s, 1H) ppm.

Example 8A

4-[(3-Amino-5-trifluoromethylphenyl)amino]-3-penten-2-one



5

1.71 g (17.0 mmol) Acetylacetone, 3.00 g (17.0 mmol) 3-amino-5-(trifluoromethyl)-phenylamine, and 0.1 g (1.6 mmol) 4-toluenesulfonic acid are dissolved in 150 ml toluene. The reaction mixture is refluxed overnight with a Dean-Stark trap to remove water. After cooling down to room temperature, the solvent is removed *in vacuo* and the residue is purified by column chromatography on silica with cyclohexane/ethylacetate mixtures as eluent. The solid obtained by column chromatography is recrystallized from cyclohexane.

10


Yield: 0.35 g (8%)

¹H-NMR (200 MHz, DMSO): δ = 2.0 (s, 3H); 2.1 (s, 3H); 5.2 (s, 1H); 5.7 (s, 2 h); 6.5 (m, 2H); 6.6 (s, 1H); 12.4 (s, 1H) ppm.

15

Preparation Examples:**Example 1**

(±)-Ethyl 5-acetyl-2-amino-4-(4-bromophenyl)-6-methyl-1-(3-nitrophenyl)-1,4-dihydro-3-pyridinecarboxylate



4.8 g (21.8 mmol) of Example 1A are dissolved in 30 ml ethanol, 4.0 g (21.8 mmol) 4-bromobenzaldehyde, 2.47 g (21.8 mmol) ethyl cyanoacetate, and 3.71 g (43.6 mmol) piperidine are added. The reaction mixture is stirred under reflux overnight. After cooling down to room temperature, the solvent is removed *in vacuo* and the residue is purified by column chromatography on silica with eluent dichloromethane.

Yield: 1.8 g (17%)

¹H-NMR (300 MHz, DMSO): δ = 1.2 (t, 3H); 1.9 (s, 3H); 2.2 (s, 3H); 4.0 (m, 2H); 4.9 (s, 1H); 6.8 (br.s, 2H); 7.3 (m, 2H); 7.5 (m, 2H); 7.8 (m, 2H); 8.2 (m, 1H); 8.4 (m, 1H) ppm.

Example 2

(-)-Ethyl 5-acetyl-2-amino-4-(4-bromophenyl)-6-methyl-1-(3-nitrophenyl)-1,4-dihydro-3-pyridinecarboxylate

- 5 The enantiomers of Example 1 are separated by chiral HPLC [silane modified poly(N-methacryloyl-L-leucin-1-menthylamide) fixed on silica; 250 x 20 mm column] with iso-hexane/ethylacetate 4:1 as eluent (20 ml/min).


$[\alpha]^{20} = -97.6^\circ$ ($\lambda = 589$ nm, methanol, $c = 456.5$ mg / 100 ml)

$^1\text{H-NMR}$ identical with Example 1.

10

Example 3

(±)-Ethyl 5-acetyl-2-amino-4-(4-bromophenyl)-6-methyl-1-[3-(trifluoromethyl)phenyl]-1,4-dihydro-3-pyridinecarboxylate



15

- 150 mg (0.62 mmol) of Example 2A are dissolved in 2 ml ethanol, 114 mg (0.62 mmol) 4-bromobenzaldehyde, 70 mg (0.62 mmol) ethyl cyanoacetate, and 105 mg (1.23 mmol) piperidine are added. The reaction mixture is stirred under reflux overnight. After cooling down to room temperature, the solvent is removed *in vacuo* and the residue is purified by column chromatography on silica with eluent dichloromethane.


20

Yield: 43 mg (13%)

¹H-NMR (300 MHz, DMSO): δ = 1.2 (t, 3H); 1.8 (s, 3H); 2.2 (s, 3H); 4.0 (m, 2H); 4.9 (s, 1H); 6.7 (br.s, 2H); 7.3 (m, 2H); 7.5 (m, 2H); 7.7 (m, 1H); 7.8 (m, 1H); 7.8 (m, 1H); 7.9 (m, 1H) ppm.

5 **Example 4**

(±)-Ethyl 5-acetyl-2-amino-4-(4-cyanophenyl)-6-methyl-1-[3-(trifluoromethyl)-phenyl]-1,4-dihydro-3-pyridinecarboxylate



10

100 mg (0.41 mmol) of Example 2A are dissolved in 2 ml ethanol, 54 mg (0.41 mmol) 4-cyanobenzaldehyde, 47 mg (0.41 mmol) ethyl cyanoacetate, and 70 mg (0.82 mmol) piperidine are added. The reaction mixture is stirred under reflux overnight. After cooling down to room temperature, the solvent is removed *in vacuo* and the residue is purified by column chromatography on silica with eluent dichloromethane.

15

Yield: 26 mg (14%)

¹H-NMR (400 MHz, DMSO): δ = 1.2 (t, 3H); 1.8 (s, 3H); 2.2 (s, 3H); 4.0 (m, 2H); 5.0 (s, 1H); 6.7 (br.s, 2H); 7.5 (m, 2H); 7.7 (m, 1H); 7.8 (m, 4H); 7.9 (m, 1H) ppm.

20

Example 5 and Example 6

(+) and (-)-Ethyl 5-acetyl-2-amino-4-(4-cyanophenyl)-6-methyl-1-[3-(trifluoro-methyl)phenyl]-1,4-dihydro-3-pyridinecarboxylate

- 5 The enantiomers of Example 4 are separated by chiral HPLC [silane modified poly(N-methacryloyl-L-leucin-1-menthylamide) fixed on silica; 250 x 20 mm column] with iso-hexane/ethylacetate 4:1 as eluent (20 ml/min).

(+)-Enantiomer (*Example 5*):

$[\alpha]^{20} = +88.4^\circ$ ($\lambda = 589$ nm, methanol, $c = 453.5$ mg / 100 ml)


- 10 (-)-Enantiomer (*Example 6*):

$[\alpha]^{20} = -91.2^\circ$ ($\lambda = 589$ nm, methanol, $c = 471.5$ mg / 100 ml)

¹H-NMRs for Example 5 and 6 identical with Example 4.

Example 7

- 15 (±)-Ethyl 5-acetyl-4-(4-bromophenyl)-6-methyl-1-[3-(trifluoromethyl)phenyl]-1,4-dihydro-3-pyridinecarboxylate




- 20 150 mg (0.62 mmol) of Example 2A are dissolved in 2 ml acetic acid, 114 mg (0.62 mmol) 4-bromobenzaldehyde, and 60 mg (0.62 mmol) ethyl propiolate are added. The reaction mixture is stirred under reflux overnight. After cooling down to room temperature, the solvent is removed *in vacuo* and the residue is purified by column chromatography on silica with eluent dichloromethane.

Yield: 56 mg (18%)

¹H-NMR (300 MHz, DMSO): δ = 1.1 (t, 3H); 1.9 (s, 3H); 2.2 (s, 3H); 4.1 (m, 2H); 5.0 (s, 1H); 7.3 (m, 3H); 7.5 (m, 2H); 7.7 (m, 2H); 7.8 (m, 1H); 7.9 (s, 1H) ppm.

5 **Example 8**

(±)-Ethyl 5-acetyl-4-(4-cyanophenyl)-6-methyl-1-[3-(trifluoromethyl)phenyl]-1,4-dihydro-3-pyridinecarboxylate



10

100 mg (0.41 mmol) of Example 2A are dissolved in 2 ml acetic acid, 54 mg (0.41 mmol) 4-cyanobenzaldehyde, and 40 mg (0.41 mmol) ethyl propiolate are added. The reaction mixture is stirred under reflux overnight. After cooling down to room temperature, the solvent is removed *in vacuo* and the residue is purified by column chromatography on silica with eluent dichloromethane.


15

Yield: 20 mg (11%)

¹H-NMR (300 MHz, DMSO): δ = 1.1 (t, 3H); 2.0 (s, 3H); 2.2 (s, 3H); 4.1 (m, 2H); 5.1 (s, 1H); 7.3 (s, 1H); 7.6 (m, 2H); 7.8 (m, 4H); 7.8 (m, 1H); 8.0 (s, 1H) ppm.

Example 9

(±)-Ethyl 5-acetyl-4-(4-cyanophenyl)-2,6-dimethyl-1-(3-methylphenyl)-1,4-dihydro-3-pyridinecarboxylate



5

100 mg (0.46 mmol) Ethyl (2E)-3-[[3-(methyl)phenyl]amino]-2-butenate, 75 mg (0.57 mmol) 4-cyanobenzaldehyde, 38 mg (0.38 mmol) 2,4-pentanedione, and 87 mg (0.76 mmol) trifluoroacetic acid are dissolved in 2 ml diisopropyl ether. The reaction mixture is stirred under reflux overnight. After cooling down to room temperature, the solvent is removed *in vacuo* and the residue is purified by preparative HPLC.


Yield: 11 mg (7%)

¹H-NMR (300 MHz, DMSO): δ = 1.2 (t, 3H); 2.0 (s, 3H); 2.0 (s, 3H); 2.2 (s, 3H); 2.4 (s, 3H); 4.1 (q, 2H); 5.1 (s, 1H); 7.0 (m, 2H); 7.3 (m, 1H); 7.4 (m, 1H); 7.5 (m, 2H); 7.8 (m, 2H) ppm.

15

Example 10

(±)-Ethyl 5-acetyl-4-(4-cyanophenyl)-2,6-dimethyl-1-[3-(trifluoromethyl)phenyl]-1,4-dihydro-3-pyridinecarboxylate



5

100 mg (0.37 mmol) of Example 3A, 60 mg (0.46 mmol) 4-cyanobenzaldehyde, 31 mg (0.31 mmol) 2,4-pentanedione, and 70 mg (0.61 mmol) trifluoroacetic acid are dissolved in 2 ml diisopropyl ether. The reaction mixture is stirred under reflux overnight. After cooling down to room temperature, the solvent is removed *in vacuo* and the residue is purified by preparative HPLC.

10


Yield: 7 mg (5%)

¹H-NMR (300 MHz, DMSO): δ = 1.2 (t, 3H); 1.9 (s, 3H); 2.0 (s, 3H); 2.3 (s, 3H); 4.1 (q, 2H); 5.1 (s, 1H); 7.5 (m, 2H); 7.6 (m, 1H); 7.7 (m, 1H); 7.8 (m, 3H); 7.9 (m, 1H) ppm.

15

Example 11

Ethyl 5-acetyl-2-amino-4-(4-(trifluoromethyl)phenyl)-6-methyl-1-[3-(trifluoromethyl)phenyl]-1,4-dihydro-3-pyridinecarboxylate



5

100 mg (0.41 mmol) of Example 2A are dissolved in 2 ml ethanol, 72 mg (0.41 mmol) 4-(trifluoromethyl)benzaldehyde, 47 mg (0.41 mmol) ethyl cyanoacetate, and 70 mg (0.82 mmol) piperidine are added. The reaction mixture is stirred under reflux overnight. After cooling down to room temperature, the solvent is removed *in vacuo* and the residue is purified by column chromatography on silica with eluent dichloromethane.


Yield: 46 mg (22%)

¹H-NMR (300 MHz, DMSO): δ = 1.2 (t, 3H); 1.9 (s, 3H); 2.2 (s, 3H); 4.0 (m, 2H); 5.0 (s, 1H); 6.8 (br. s, 2H); 7.5 (m, 2H); 7.7 (m, 3H); 7.7 (m, 1H); 7.8 (m, 1H); 7.9 (m, 1H) ppm.

15

Example 12

Ethyl 5-acetyl-2-amino-4-(4-cyanophenyl)-6-methyl-1-[3-methylphenyl]-1,4-dihydro-3-pyridinecarboxylate



5

100 mg (0.53 mmol) of Example 4A are dissolved in 2 ml ethanol, 69 mg (0.53 mmol) 4-cyanobenzaldehyde, 60 mg (0.53 mmol) ethyl cyanoacetate, and 90 mg (1.06 mmol) piperidine are added. The reaction mixture is stirred under reflux overnight. After cooling down to room temperature, the solvent is removed *in vacuo* and the residue is purified by column chromatography on silica with eluent dichloromethane.


Yield: 21 mg (10%)

¹H-NMR (200 MHz, DMSO): δ = 1.2 (t, 3H); 1.9 (s, 3H); 2.2 (s, 3H); 4.0 (m, 2H); 5.0 (s, 1H); 6.7 (br. s, 2H); 7.2 (m, 2H); 7.3 (m, 1H); 7.5 (m, 3H); 7.8 (m, 2H) ppm.

15

Example 13

5-Acetyl-2-amino-4-(4-cyanophenyl)-6-methyl-1-[3-(trifluoromethyl)phenyl]-1,4-dihydro-3-pyridinecarboxamide



5

750 mg (3.08 mmol) of Example 2A are dissolved in 5 ml ethanol, 404 mg (3.08 mmol) 4-cyanobenzaldehyde, 260 mg (3.08 mmol) cyanoacetamide, and 26 mg (0.31 mmol) piperidine are added. The reaction mixture is stirred under reflux overnight. After cooling down to room temperature, the solvent is removed *in vacuo* and the residue is purified by column chromatography on silica with eluent dichloromethane.

10


Yield: 160 mg (12%)

¹H-NMR (300 MHz, DMSO): δ = 1.8 (s, 3H); 2.2 (s, 3H); 4.9 (s, 1H); 6.7 (br. s, 2H); 6.9 (br. s, 2H); 7.5 (m, 3H); 7.8 (m, 2H); 7.9 (m, 1H); 8.0 (m, 2H) ppm.

15

Example 14

5-Acetyl-4-(4-cyanophenyl)-2-imino-N,N,6-trimethyl-1-[3-(trifluoromethyl)phenyl]-1,2,3,4-tetrahydro-3-pyridinecarboxamide



5

750 mg (3.08 mmol) of Example 2A are dissolved in 5 ml ethanol, 404 mg (3.08 mmol) 4-cyanobenzaldehyde, 260 mg (3.08 mmol) 2-cyano-N,N-dimethylacetamide, and 26 mg (0.31 mmol) piperidine are added. The reaction mixture is stirred under reflux overnight. After cooling down to room temperature, the solvent is removed *in vacuo* and the residue is purified by column chromatography on silica with eluent dichloromethane.

10


Yield: 88 mg (6%)

¹H-NMR (300 MHz, DMSO): δ = 2.0 (s, 3H); 2.1 (s, 3H), 2.5 (s, 3H); 2.9 (s, 3H); 4.1 (d, 1H); 4.5 (d, 1H); 7.6 (m, 3H); 7.7 (m, 1H); 7.8 (m, 3H); 8.2 (s, 1H) ppm.

15

Example 15

5-Acetyl-2-amino-4-(4-cyanophenyl)-6-methyl-1-[3-(trifluoromethyl)phenyl]-1,4-dihydro-3-pyridinecarbonitrile



5

750 mg (3.08 mmol) of Example 2A are dissolved in 5 ml ethanol, 404 mg (3.08 mmol) 4-cyanobenzaldehyde, 204 mg (3.08 mmol) malononitrile, and 26 mg (0.31 mmol) piperidine are added. The reaction mixture is stirred under reflux overnight. After cooling down to room temperature, the solvent is removed *in vacuo* and the residue is purified by column chromatography on silica with eluent cyclohexane/ethyl acetate.

10


Yield: 43 mg (3%)

LC-MS: retention time 3.17 min., $m/z = 423$ $[M+H]^+$.

15

Example 16

Ethyl 5-acetyl-2-amino-4-(4-cyanophenyl)-1-[3-iodo-5-(trifluoromethyl)phenyl]-6-methyl-1,4-dihydro-3-pyridinecarboxylate



5


200 mg (0.54 mmol) of Example 5A are dissolved in 2.5 ml ethanol, 71.1 mg (0.54 mmol) 4-cyanobenzaldehyde, 61.3 mg (0.54 mmol) ethyl cyanoacetate, and 4.6 mg (0.05 mmol) piperidine are added. The reaction mixture is stirred under reflux for 30 h. After cooling down to room temperature, the solvent is removed *in vacuo* and the residue is purified by column chromatography on silica with eluent cyclohexane/ethyl acetate. The solid obtained by column chromatography is dissolved in ethanol. Water is added until a solid precipitates. The suspension is filtered and the solvent of the filtrate is removed *in vacuo*.

15 Yield: 18 mg (5%)

¹H-NMR (300 MHz, DMSO): δ = 1.1 (t, 3H); 1.8 (s, 3H); 2.2 (s, 3H); 4.0 (q, 2H); 4.9 (s, 1H); 6.9 (s, 2H); 7.5 (d, 2H); 7.7 (d, 2H); 7.8 (s, 1H); 8.1 (s, 1H); 8.2 (s, 1H) ppm.

Example 17

Ethyl 5-acetyl-2-amino-4-(4-cyanophenyl)-1-{3-[(diethylamino)methyl]phenyl}-6-methyl-1,4-dihydro-3-pyridinecarboxylate



5

500 mg (1.92 mmol) of Example 6A are dissolved in 3.0 ml ethanol, 265 mg (1.92 mmol) 4-cyanobenzaldehyde, 217 mg (1.92 mmol) ethyl cyanoacetate, and 49 mg (0.58 mmol) piperidine are added. The reaction mixture is stirred under reflux for 48 h. 115 mg (1.34 mmol) piperidine are added and the mixture is stirred under reflux for another 6 h. After cooling down to room temperature, the solvent is removed *in vacuo* and the residue is purified by preparative HPLC with eluent acetonitrile/water.


Yield: 57 mg (5%)

¹H-NMR (200 MHz, DMSO): δ = 1.0 (t, 6H); 1.2 (t, 3H); 1.9 (s, 3H); 2.2 (s, 3H); 2.5 (m, 4H); 3.6 (s, 2H); 4.0 (q, 2H); 5.0 (s, 1H); 6.7 (br. s, 2H); 7.2 (m, 2H); 7.5 (m, 4H); 7.8 (d, 2H) ppm.

15

Example 18

Ethyl 5-acetyl-2-amino-4-(4-cyanophenyl)-1-[3-methoxy-5-(trifluoromethyl)phenyl]-6-methyl-1,4-dihydro-3-pyridinecarboxylate



5

200 mg (0.73 mmol) of Example 7A are dissolved in 2.5 ml ethanol, 95.9 mg (0.73 mmol) 4-cyanobenzaldehyde, 82.8 mg (0.73 mmol) ethyl cyanoacetate, and 6.2 mg (0.07 mmol) piperidine are added. The reaction mixture is stirred under reflux overnight. After cooling down to room temperature, the solvent is removed *in vacuo* and the residue is purified by preparative HPLC with eluent acetonitrile/water.

10

Yield: 59 mg (15%)


¹H-NMR (300 MHz, DMSO): δ = 1.2 (t, 3H); 1.9 (s, 3H); 2.2 (s, 3H); 3.9 (s, 3H); 4.0 (q, 2H); 5.0 (s, 1H); 6.8 (br. s, 2H); 7.3 (m, 2H); 7.4 (s, 1H); 7.5 (d, 2H); 7.8 (d, 2H)

15

ppm.

Example 19

Ethyl 5-acetyl-2-amino-1-[3-amino-5-(trifluoromethyl)phenyl]-4-(4-cyanophenyl)-6-methyl-1,4-dihydro-3-pyridinecarboxylate




200 mg (0.77 mmol) of Example 8A are dissolved in 2.5 ml ethanol, 107 mg (0.77 mmol) 4-cyanobenzaldehyde, 87.6 mg (0.77 mmol) ethyl cyanoacetate, and 20 mg (0.23 mmol) piperidine are added. The reaction mixture is stirred under reflux overnight. After cooling down to room temperature, the solvent is removed *in vacuo* and the residue is purified by preparative HPLC with eluent acetonitrile/water.

Yield: 50.5 mg (12%)

¹H-NMR (300 MHz, DMSO): δ = 1.2 (t, 3H); 2.0 (s, 3H); 2.3 (s, 3H); 4.1 (q, 2H); 5.0 (s, 1H); 6.0 (s, 2H); 6.7 (s, 2H); 6.8 (s, 2H); 7.1 (s, 1H); 7.4 (d, 2H); 7.7 (d, 2H) ppm.

Example 20

Ethyl 5'-acetyl-2'-amino-6'-methyl-1'-[3-(trifluoromethyl)phenyl]-1',4'-dihydro-3,4'-bipyridine-3'-carboxylate



5

100 mg (0.41 mmol) of Example 2A are dissolved in 5 ml ethanol, 44 mg (0.41 mmol) nicotinaldehyde, 47 mg (0.41 mmol) ethyl cyanoacetate, and 70 mg (0.82 mmol) piperidine are added. The reaction mixture is stirred under reflux overnight. After cooling down to room temperature, the solvent is removed *in vacuo* and the residue is purified by column chromatography on silica with eluent dichloromethane.

Yield: 19 mg (10%)
¹H-NMR (300 MHz, DMSO): δ = 2.0 (s, 3H); 1.8 (s, 3H); 2.2 (s, 3H); 4.0 (m, 2H);
15 4.9 (s, 1H); 6.7 (m, 2H); 7.3 (m, 1H); 7.7 (m, 2H); 7.8 (m, 2H); 7.9 (m, 1H); 8.4 (m, 1H); 8.5 (m, 1H) ppm.

C. Operative examples relating to pharmaceutical compositions

The compounds according to the invention can be converted into pharmaceutical preparations as follows:

5

Tablet:**Composition:**

100 mg of the compound of Example 1, 50 mg of lactose (monohydrate), 50 mg of maize starch (native), 10 mg of polyvinylpyrrolidone (PVP 25) (from BASF, Ludwigshafen, Germany) and 2 mg of magnesium stearate.

Tablet weight 212 mg, diameter 8 mm, curvature radius 12 mm.

Preparation:

The mixture of active component, lactose and starch is granulated with a 5% solution (m/m) of the PVP in water. After drying, the granules are mixed with magnesium stearate for 5 min. This mixture is moulded using a customary tablet press (tablet format, see above). The moulding force applied is typically 15 kN.

15

Orally administrable suspension:**Composition:**

1000 mg of the compound of Example 1, 1000 mg of ethanol (96%), 400 mg of Rhodigel (xanthan gum from FMC, Pennsylvania, USA) and 99 g of water.

A single dose of 100 mg of the compound according to the invention is provided by 10 ml of oral suspension.

20


Preparation:

The Rhodigel is suspended in ethanol and the active component is added to the suspension. The water is added with stirring. Stirring is continued for about 6h until the swelling of the Rhodigel is complete.

25

We claim

1. Compounds of the general formula (I)



wherein

R^1 , R^2 , R^3 , R^8 , R^9 and R^{10} independently from each other represent hydrogen, halogen, nitro, cyano, trifluoromethyl, C_1 - C_6 -alkyl, C_1 - C_6 -alkanoyl, hydroxy, C_1 - C_6 -alkoxy, trifluoromethoxy, amino, mono- or di- C_1 - C_6 -alkylamino, C_1 - C_6 -acylamino, C_1 - C_6 -alkoxycarbonylamino, carboxyl, C_1 - C_6 -alkoxycarbonyl or phenyl, wherein C_1 - C_6 -alkyl, C_1 - C_6 -alkanoyl, C_1 - C_6 -alkoxy, mono- or di- C_1 - C_6 -alkylamino and C_1 - C_6 -acylamino can be further substituted with one to three identical or different radicals selected from the group consisting of hydroxy, C_1 - C_4 -alkoxy, amino, mono- and di- C_1 - C_4 -alkylamino,


R^4 represents C_1 - C_6 -alkyl, trifluoromethyl or phenyl,

R^5 represents C_1 - C_4 -alkyl, which can be substituted with one to three identical or different radicals selected from the group consisting of hydroxy, C_1 - C_4 -alkoxy, amino, mono- and di- C_1 - C_4 -alkylamino,

R^6 represents cyano, aminocarbonyl, mono- and di- C_1 - C_4 -alkylamino-carbonyl, carboxyl or C_1 - C_6 -alkoxycarbonyl, wherein the alkoxy moiety can be further substituted with a radical selected from the group consisting of hydroxy, C_1 - C_4 -alkoxy, amino, mono- and di- C_1 - C_4 -alkylamino,

or

R^6 represents a moiety of the formula



or

wherein R^A is selected from the group consisting of hydrogen and C_1 - C_6 -alkyl,

R^7 represents hydrogen, C_1 - C_4 -alkyl or amino,

X^1 , X^2 , X^3 and X^4 independently from each other represent CH or N, wherein ring A contains either 0, 1 or 2 nitrogen atoms,

and

Y^1 , Y^2 , Y^3 , Y^4 and Y^5 independently from each other represent CH or N, wherein ring B contains either 0, 1 or 2 nitrogen atoms and their salts, hydrates and/or solvates.

2. Compounds of general formula (I) according to claim 1, wherein

R¹ represents hydrogen,

5 R², R³, R⁹ and R¹⁰ independently from each other represent hydrogen, halogen, nitro, cyano, trifluoromethyl, C₁-C₄-alkyl, C₁-C₄-alkanoyl, hydroxy, C₁-C₄-alkoxy, trifluoromethoxy, amino, C₁-C₄-dialkylamino, C₁-C₄-acylamino, methoxy-carbonylamino, tert.-butoxycarbonyl-
10 amino, carboxyl, methoxycarbonyl or ethoxycarbonyl, wherein C₁-C₄-alkyl, C₁-C₄-alkanoyl, C₁-C₄-alkoxy, C₁-C₄-dialkylamino and C₁-C₄-acylamino can be further substituted with one to two identical or different radicals selected from the group consisting of hydroxy, methoxy, ethoxy, amino, dimethylamino and diethylamino,

15 R⁴ represents methyl, ethyl, trifluoromethyl or phenyl,

R⁵ represents methyl or ethyl,

20 R⁶ represents cyano, aminocarbonyl, methylaminocarbonyl, dimethylaminocarbonyl, carboxyl or C₁-C₄-alkoxycarbonyl, wherein the alkoxy moiety can be further substituted with a radical selected from the group consisting of hydroxy, methoxy, ethoxy, amino, mono- and di-C₁-C₄-alkylamino,

25 R⁷ represents hydrogen, methyl, ethyl or amino,

R⁸ represents hydrogen,

X¹, X² and X³ represent CH,

30

X⁴ represents CH or N,

and

Y^1, Y^2, Y^3, Y^4 and Y^5 represent CH.

5

3. Compounds of general formula (I) according to claim 1 or 2, wherein

R^1 and R^2 represent hydrogen,

10 R^3 represents fluoro, chloro, bromo, nitro, cyano, trifluoromethyl, methyl, methoxy or hydroxy,

R^4 represents methyl or trifluoromethyl,

15 R^5 represents methyl,

R^6 represents cyano, aminocarbonyl, methylaminocarbonyl, dimethylaminocarbonyl, carboxyl, methoxycarbonyl or ethoxycarbonyl,

20 R^7 represents hydrogen, methyl or amino,

R^8 and R^9 represent hydrogen,

25 R^{10} represents fluoro, chloro, bromo, nitro, cyano, trifluoromethyl or methyl,

X^1, X^2, X^3 and X^4 represent CH,


and

30

Y^1, Y^2, Y^3, Y^4 and Y^5 represent CH.


4. Compounds of general formula (I) according to claim 1, 2 or 3, wherein R³ is cyano, which is located in para-position relative to the 1,4-dihydropyridine ring.
5. Compounds of general formula (I) according to at least one of claims 1 to 4, wherein R⁴ is methyl.
6. Compounds of general formula (I) according to at least one of claims 1 to 5, wherein R⁵ is methyl.
7. Compounds of general formula (I) according to at least one of claims 1 to 6, wherein R⁶ is methoxycarbonyl or ethoxycarbonyl.
8. Compounds of general formula (I) according to at least one of claims 1 to 7, wherein R⁷ is hydrogen, methyl or amino.
9. Compounds of general formula (I) according to at least one of claims 1 to 8, wherein R¹⁰ is trifluoromethyl, which is attached to Y².
10. Processes for synthesizing the compounds of general formula (I) as defined in claims 1 to 9, characterized in that either

[A] compounds of the general formula (II)



wherein R^4 , R^5 , R^8 , R^9 , R^{10} and Y^1 to Y^5 have the meaning described in claim 1,

are condensed in the presence of a base with compounds of the general formulas (III) and (IV)



(III)




(IV),

wherein R^1 , R^2 , R^3 and X^1 to X^4 have the meaning described in claim 1,

and

R^{11} represents cyano or C_1 - C_6 -alkoxycarbonyl,


to give compounds of the general formula (Ia)



(Ia),

or


[B] compounds of the general formulas (II) and (III) are condensed in the presence of an acid with compounds of the general formula (V)



wherein R^{12} represents $\text{C}_1\text{-C}_6\text{-alkyl}$,

5


to give compounds of the general formula (Ib)



or

10

[C] compounds of the general formula (VI)




15

wherein $\text{R}^8, \text{R}^9, \text{R}^{10}, \text{R}^{12}$ and Y^1 to Y^5 have the meaning described in claim 1,

and


R^{13} represents C_1 - C_4 -alkyl,

are condensed in the presence of an acid or a base with compounds of the
 5 general formulas (III) and (VII)



wherein R^4 and R^5 have the meaning described in claim 1,

10 to give compounds of the general formula (Ic)



- 15 11. The composition containing at least one compound of general formula (I) as defined in claims 1 to 9 and a pharmacologically acceptable diluent.
12. A composition according to claim 11 for the treatment of acute and chronic inflammatory processes.

13. The process for the preparation of compositions according to claim 11 and 12 characterized in that the compounds of general formula (I) as defined in claims 1 to 9 together with customary auxiliaries are brought into a suitable application form.
- 5
14. Use compounds of general formula (I) as defined in claims 1 to 9 for the preparation of medicaments.
15. Use according to claim 14 for the preparation of medicaments for the treatment of acute and chronic inflammatory processes.
- 10
16. Use according to claim 15, wherein the process is chronic obstructive pulmonary disease.
- 15
17. Process for controlling chronic obstructive pulmonary disease in humans and animals by administration of a neutrophil elastase inhibitory amount of at least one compound according to any of Claims 1 to 9.

International Application No.

A. CLASSIFICATION OF SUBJECT MATTER

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C07D A61K A61P

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, CHEM ABS Data, WPI Data, PAJ, BIOSIS, EMBASE

C. DOCUMENTS CONSIDERED TO BE RELEVANT

X

Further documents are listed in the continuation of box C.



Patent family members are listed in annex.

* Special categories of cited documents :

- *T later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

- *X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- *Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- *G* document member of the same patent family

Date of the actual completion of the international search

19 March 2003

Date of mailing of the international search report

04/04/2003

Name and mailing address of the ISA
European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax (+31-70) 340-3016

Authorized officer

Frelon, D

INTERNATIONAL SEARCH REPORT

International Application No

PCT/EP 02/13931

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	DATABASE CA 'Online! CHEMICAL ABSTRACTS SERVICE, COLUMBUS, OHIO, US; SEKI, TAKEJI ET AL: "Preparation of 1,4-dihydropyridine compounds as antiallergics and antiinflammatories" retrieved from STN Database accession no. 124:343126 XP002235303 abstract & JP 08 040904 A (NIKKEN CHEMICALS CO LTD, JAPAN) 13 February 1996 (1996-02-13) _____	1-17

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/EP 02/13931

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
WO 9505823	A	02-03-1995	DE 4328884 A1	02-03-1995
			AU 7613594 A	21-03-1995
			BG 100367 A	31-07-1996
			CA 2170285 A1	02-03-1995
			CN 1129906 A	28-08-1996
			CZ 9600438 A3	15-05-1996
			WO 9505823 A1	02-03-1995
			EP 0714301 A1	05-06-1996
			HU 75302 A2	28-05-1997
			JP 9501923 T	25-02-1997
			NO 960777 A	10-04-1996
			PL 313116 A1	10-06-1996
FR 2528425	A	16-12-1983	DE 3222367 A1	15-12-1983
			BE 897035 A1	13-12-1983
			CA 1228549 A1	27-10-1987
			ES 8403456 A1	16-06-1984
			FR 2528425 A1	16-12-1983
			IT 1175920 B	12-08-1987
			JP 1840675 C	25-04-1994
			JP 5053774 B	10-08-1993
			JP 59005114 A	12-01-1984
			ZA 8304344 A	28-03-1984
EP 0451654	A	16-10-1991	DE 4011695 A1	17-10-1991
			AT 111079 T	15-09-1994
			AU 642992 B2	04-11-1993
			AU 7425391 A	17-10-1991
			CA 2040062 A1	12-10-1991
			DE 59102788 D1	13-10-1994
			DK 451654 T3	13-02-1995
			EP 0451654 A2	16-10-1991
			ES 2063998 T3	16-01-1995
			FI 911700 A	12-10-1991
			HU 59905 A2	28-07-1992
			HU 71485 A2	28-11-1995
			HU 9500334 A3	30-10-1995
			IE 911200 A1	23-10-1991
			IL 97796 A	15-03-1995
			JP 3012352 B2	21-02-2000
			JP 4234847 A	24-08-1992
			KR 180741 B1	20-03-1999
			NZ 237722 A	26-10-1995
			NZ 250085 A	24-03-1997
			PT 97300 A ,B	31-01-1992
			US 5432185 A	11-07-1995
			US 5234935 A	10-08-1993
			ZA 9102652 A	29-01-1992
JP 8040904	A	13-02-1996	JP 3337829 B2	28-10-2002